

NEW BEAUVILLE SURFACES AND FINITE SIMPLE GROUPS

SHELLY GARION, MATTEO PENEGINI

ABSTRACT. In this paper we construct new Beauville surfaces with group either $\mathrm{PSL}(2, p^e)$, or belonging to some other families of finite simple groups of Lie type of low Lie rank, or an alternating group, or a symmetric group, proving a conjecture of Bauer, Catanese and Grunewald. The proofs rely on probabilistic group theoretical results of Liebeck and Shalev, on classical results of Macbeath and on recent results of Marion.

1. INTRODUCTION

A *Beauville surface* S (over $\mathbb{C}$) is a particular kind of surface isogenous to a higher product of curves, i.e., $S = (C_1 \times C_2)/G$ is a quotient of a product of two smooth curves C_1, C_2 of genera at least two, modulo a free action of a finite group G , which acts faithfully on each curve. For Beauville surfaces the quotients C_i/G are isomorphic to $\mathbb{P}^1$ and both projections $C_i \rightarrow C_i/G \cong \mathbb{P}^1$ are coverings branched over three points. A Beauville surface is in particular a minimal surface of general type.

Beauville surfaces were introduced by F. Catanese in [Cat00], inspired by a construction of A. Beauville (see [B]). The two authors were interested in finding new examples of surfaces with $p_g = q = 0$ and of general type, which provide an interesting class of surfaces (see e.g. [BCG08]). As a matter of fact a Beauville surface has $q = 0$, but p_g can attain any non negative value. Since [Cat00] had appeared, many authors have been studying Beauville surfaces, see [BCG05, BCG06, BCG08, FG, FGJ, FJ].

Working out the definition of Beauville surfaces one sees that there is a pure group theoretical condition which characterizes the groups of Beauville surfaces: the existence of what in [BCG05] and [BCG06] is called a "Beauville structure".

Definition 1.1. *An unmixed Beauville structure for a finite group G is a quadruple $(x_1, y_1; x_2, y_2)$ of elements of G , which determines two triples $T_i := (x_i, y_i, z_i)$ ($i = 1, 2$) of elements of G such that :*

- (i) $x_i y_i z_i = 1$,
- (ii) $\langle x_i, y_i \rangle = G$,
- (iii) $\Sigma(T_1) \cap \Sigma(T_2) = \{1\}$, where

$$\Sigma(T_i) := \bigcup_{g \in G} \bigcup_{j=1}^{\infty} \{g x_i^j g^{-1}, g y_i^j g^{-1}, g z_i^j g^{-1}\}.$$

Moreover, $\tau_i := (\text{ord}(x_i), \text{ord}(y_i), \text{ord}(z_i))$ is called the *type* of T_i , and a type which satisfies the condition:

$$\frac{1}{\text{ord}(x_i)} + \frac{1}{\text{ord}(y_i)} + \frac{1}{\text{ord}(z_i)} < 1$$

is called *hyperbolic*.

A detailed explanation of this link between surfaces and finite groups will be given in Section 2, where we shall present a more general definition related to surfaces isogenous to a higher product.

Therefore, the question of which finite groups G admit an unmixed Beauville structure was raised. The following Theorem summarizes the known results.

Theorem 1.2. *The following groups admit an unmixed Beauville structure:*

- (1) *The alternating groups A_n admit unmixed Beauville structures if and only if $n \geq 6$;*
- (2) *The symmetric groups S_n admit unmixed Beauville structures if and only if $n \geq 5$;*
- (3) *The groups $\text{SL}(2, p)$ and $\text{PSL}(2, p)$ for every prime $p \neq 2, 3, 5$;*
- (4) *The Suzuki groups $\text{Sz}(2^p)$, where p is an odd prime;*
- (5) *A finite abelian group G admits an unmixed Beauville structure if and only if $G = (\mathbb{Z}/n\mathbb{Z})^2$ with $(n, 6) = 1$;*
- (6) *For every prime p , there exists a p -group which admits an unmixed Beauville structure.*

Proof. Part (1) was proven in [BCG05], [BCG06] for n large enough, and it was later generalized in [FG]. Part (2) was proven for $n \geq 7$ in [BCG06], and it was later improved in [FG]. Parts (3), (4) and (5) appeared in [BCG05] (for part (5) see also [Cat00]). Part (6) is a consequence of (5) for $p \geq 5$, and the proof for $p = 2, 3$ appeared in [FGJ]. $\square$

The question of which finite groups admit an unmixed Beauville structure is deeply related to the question of which finite groups are quotients of certain triangle groups, which was widely investigated (see [Co90, Co10] for a survey). Indeed, conditions (i) and (ii) of Definition 1.1 clearly imply that two certain triangle groups surject onto the finite group G . However, the question about Beauville structures is somewhat more delicate, due to condition (iii) of Definition 1.1.

The following Theorem regarding alternating groups, generalizing part (1) of Theorem 1.2, was conjectured by Bauer, Catanese and Grunewald in [BCG05, BCG06], who were inspired by the proof of Everitt [Ev] to Higman's Conjecture that every hyperbolic triangle group surjects to all but finitely many alternating groups.

Theorem 1.3. *Let $(r_1, s_1, t_1), (r_2, s_2, t_2)$ be two hyperbolic types. Then almost all alternating groups A_n admit an unmixed Beauville structure $(x_1, y_1; x_2, y_2)$ where $(x_1, y_1, (x_1 y_1)^{-1})$ has type (r_1, s_1, t_1) and $(x_2, y_2, (x_2 y_2)^{-1})$ has type (r_2, s_2, t_2) .*

A similar Theorem also applies for symmetric groups.

Theorem 1.4. *Let $(r_1, s_1, t_1), (r_2, s_2, t_2)$ be two hyperbolic types, and assume that at least two of (r_1, s_1, t_1) are even and at least two of (r_2, s_2, t_2) are even.*

Then almost all symmetric groups S_n admit an unmixed Beauville structure $(x_1, y_1; x_2, y_2)$ where $(x_1, y_1, (x_1 y_1)^{-1})$ has type (r_1, s_1, t_1) and $(x_2, y_2, (x_2 y_2)^{-1})$ has type (r_2, s_2, t_2) .

The proofs of both Theorems are presented in Section 3.1, and are based on results of Liebeck and Shalev [LS04], who gave an alternative proof, based on probabilistic group theory, to Higman's Conjecture. We also provide similar Theorems for surfaces isogenous to a higher product not necessarily Beauville.

The following Theorem generalizes part (3) of Theorem 1.2.

Theorem 1.5. *Let p be a prime number, and assume that $q = p^e$ is at least 7. Then the group $\mathrm{PSL}(2, q)$ admits an unmixed Beauville structure.*

Its following Corollary is analogous to Theorem 1.3 for the family of groups $\{\mathrm{PSL}(2, p)\}_{p \text{ prime}}$.

Corollary 1.6. *Let $r, s > 5$ be two relatively prime integers. Then there are infinitely many primes p for which the group $\mathrm{PSL}(2, p)$ admits an unmixed Beauville structure $(x_1, y_1; x_2, y_2)$ where $(x_1, y_1, (x_1 y_1)^{-1})$ has type (r, r, r) and $(x_2, y_2, (x_2 y_2)^{-1})$ has type (s, s, s) .*

This Theorem and its Corollary are proved in Section 3.2. The proof is based on properties of the groups $\mathrm{PSL}(2, q)$ and on results of Macbeath [Ma].

Moreover, one can generalize Theorem 1.5, as well as part (4) of Theorem 1.2, and prove similar results regarding some other families of finite simple groups of Lie type of low Lie rank, provided that their defining field is large enough.

Theorem 1.7. *The following finite simple groups of Lie type $G = G(q)$ admit an unmixed Beauville structure, provided that q is large enough.*

- (1) *Suzuki groups, $G = \mathrm{Sz}(q) = {}^2 B_2(q)$, where $q = 2^{2e+1}$;*
- (2) *Ree groups, $G = {}^2 G_2(q)$, where $q = 3^{2e+1}$;*
- (3) *$G = G_2(q)$, where $q = p^e$ for some prime number $p > 3$;*
- (4) *$G = {}^3 D_4(q)$, where $q = p^e$ for some prime number $p > 3$;*
- (5) *$G = \mathrm{PSL}(3, q)$, where $q = p^e$ for some prime p ;*
- (6) *$G = \mathrm{PSU}(3, q)$, where $q = p^e$ for some prime p .*

This Theorem is proved in Section 3.3, and the proof is based on recent probabilistic group theoretical results of Marion [Mar3.09, Mar9.09], who investigated the possible surjection of certain triangle groups onto finite simple groups of Lie type of low Lie rank.

Another Conjecture of Bauer, Catanese and Grunewald [BCG05, BCG06] is that all finite simple non-abelian groups, except A_5 , admit an unmixed Beauville structure. Indeed, Theorems 1.5 and 1.7 are a first step towards a possible proof of this Conjecture for all finite simple groups of Lie type (assuming that their defining field is large enough). Moreover, in the same direction of Theorems 1.3 and 1.4, and inspired by conjectures of Liebeck and Shalev [LS05] (see also Section 3.3.2), we propose the following Conjecture.

Conjecture 1.8. *Let $(r_1, s_1, t_1), (r_2, s_2, t_2)$ be two hyperbolic types. If G is a finite simple classical group of Lie type of Lie rank large enough, then it admits an unmixed Beauville structure $(x_1, y_1; x_2, y_2)$, where $(x_1, y_1, (x_1 y_1)^{-1})$ has type (r_1, s_1, t_1) and $(x_2, y_2, (x_2 y_2)^{-1})$ has type (r_2, s_2, t_2) .*

This paper is organized as follows. In Section 2 we shall present the geometrical background and explain the link between geometry and group theory. In Section 3 one can find the proofs of the main results.

Remark 1.9. After completing this manuscript, it was brought to our attention that Fuertes and Jones [FJ], have independently and simultaneously constructed unmixed Beauville structures for the groups $\mathrm{PSL}(2, q)$, the Suzuki groups $G = \mathrm{Sz}(q) = {}^2B_2(q)$ and the Ree groups $G = {}^2G_2(q)$, thus proving some of our results appearing in Theorems 1.5 and 1.7. However, their constructions are of different type.

Following the posting of these results on the arXiv, many authors have intensively investigated Beauville surfaces and due to [GLL], [GM] and [FMP] the Conjecture of Bauer, Catanese and Grunewald, that all finite simple non-abelian groups, except A_5 , admit an unmixed Beauville structure, is now proven.

2. GEOMETRICAL BACKGROUND ON RAMIFICATION STRUCTURES

We shall denote by S a smooth irreducible complex projective surface of general type. We shall also use the standard notation in surface theory, hence we denote by $p_g := h^0(S, \Omega_S^2)$ the *geometric genus* of S , $q := h^0(S, \Omega_S^1)$ the *irregularity* of S , $\chi(S) = 1 + p_g - q$ the *holomorphic Euler-Poincaré characteristic*, $e(S)$ the *topological Euler number*, and K_S^2 the *self-intersection of the canonical divisor* (see e.g. [BHPV]). In this section, C will always denote a smooth compact complex curve and $g(C)$ will be its *genus*.

Definition 2.1. A surface S is said to be isogenous to a higher product of curves if and only if S is a quotient $(C_1 \times C_2)/G$, where C_1 and C_2 are curves of genus at least two, and G is a finite group acting freely on $C_1 \times C_2$.

In [Cat00] it is proven that any surface isogenous to a higher product has a unique minimal realization as a quotient $(C_1 \times C_2)/G$, where G is a finite group acting freely and with the property that no element acts trivially on one of the two factors C_i . From now on we shall work only with minimal realization.

We have two cases: the *mixed* case where the action of G exchanges the two factors (and then C_1 and C_2 are isomorphic), and the *unmixed* case where G acts diagonally on their product.

A surface S isogenous to a higher product is in particular a minimal surface of general type and it has

$$K_S^2 = 8\chi(S), \quad 4\chi(S) = e(S), \quad \text{and}$$

$$(1) \quad \chi(S) = \frac{(g(C_1) - 1)(g(C_2) - 1)}{|G|},$$

by Theorem 3.4 of [Cat00]. Moreover, by Serrano [S, Proposition 2.2],

$$(2) \quad q(S) = g(C_1/G) + g(C_2/G),$$

see also [Cat00] paragraph 3.

A special case of surfaces isogenous to a higher product is given by Beauville surfaces, which were also defined in [Cat00].

Definition 2.2. A Beauville surface is a surface isogenous to a higher product $S = (C_1 \times C_2)/G$, which is rigid, i.e., it has no nontrivial deformation.

Remark 2.3. Every Beauville surface of mixed type has an unramified double covering which is a Beauville surface of unmixed type.

The rigidity property of the Beauville surfaces is equivalent to the fact that $C_i/G \cong \mathbb{P}^1$ and that the projections $C_i \rightarrow C_i/G \cong \mathbb{P}^1$ are branched in three points, for $i = 1, 2$. Moreover, by Equation (2) one has $q(S) = 0$.

In the following we shall consider only the *unmixed* case.

From the above Remark one can see that studying Beauville surfaces (as well as surfaces isogenous to a higher product in general) is strictly linked to the study of branched covering of complex curves. We shall recall Riemann's Existence Theorem which translates the geometric problem of constructing branch covering into a group theoretical problem. We state it first in great generality.

Definition 2.4. Let $g', m_1, \dots, m_r$ be positive integers. An orbifold surface group of type $(g' \mid m_1, \dots, m_r)$ is a group presented as follows:

$$\Gamma(g' \mid m_1, \dots, m_r) := \langle a_1, b_1, \dots, a_{g'}, b_{g'}, c_1, \dots, c_r \mid c_1^{m_1} = \dots = c_r^{m_r} = \prod_{k=1}^{g'} [a_k, b_k] c_1 \cdot \dots \cdot c_r = 1 \rangle.$$

If $g' = 0$ it is called a polygonal group, if $g' = 0$ and $r = 3$ it is called a triangle group.

We remark that an orbifold surface group is in particular a *Fuchsian group* (see e.g. [LS04]).

The following is a reformulation of *Riemann's Existence Theorem*:

Theorem 2.5. A finite group G acts as a group of automorphisms on some compact Riemann surface C of genus g if and only if there are natural numbers $g', m_1, \dots, m_r$, and an orbifold homomorphism

$$\theta: \Gamma(g' \mid m_1, \dots, m_r) \rightarrow G$$

such that $\text{ord}(\theta(c_i)) = m_i \ \forall i$ and such that the Riemann - Hurwitz relation holds:

$$(3) \quad 2g - 2 = |G| \left(2g' - 2 + \sum_{i=1}^r \left(1 - \frac{1}{m_i} \right) \right).$$

If this is the case, then g' is the genus of $C' := C/G$. The G -cover $C \rightarrow C'$ is branched in r points $p_1, \dots, p_r$ with branching indices $m_1, \dots, m_r$, respectively.

Moreover, if we denote by $x_i \in G$ the image of c_i under θ , then

$$\Sigma(x_1, \dots, x_r) := \cup_{a \in G} \cup_{i=0}^{\infty} \{ax_1^i a^{-1}, \dots, ax_r^i a^{-1}\},$$

is the set of stabilizers for the action of G on C .

If we restrict ourselves to the case where the quotient curve is isomorphic to $\mathbb{P}^1$ then the Theorem suggests the following definition.

Definition 2.6. Let G be a finite group and $r \in \mathbb{N}$ with $r \geq 2$.

- An r -tuple $T = (x_1, \dots, x_r)$ of elements of G is called a spherical r -system of generators of G if $\langle x_1, \dots, x_r \rangle = G$ and $x_1 \cdot \dots \cdot x_r = 1$.
- We say that T is of type $\tau := (m_1, \dots, m_r)$ if the orders of $(x_1, \dots, x_r)$ are respectively $(m_1, \dots, m_r)$.
- We shall denote:

$$\mathcal{S}(G, \tau) := \{\text{spherical } r\text{-systems for } G \text{ of type } \tau\}.$$

- Moreover, two spherical r_i -systems $T_1 = (x_1, \dots, x_{r_1})$ and $T_2 = (x_1, \dots, x_{r_2})$ are said to be disjoint, if:

$$(4) \quad \Sigma(T_1) \cap \Sigma(T_2) = \{1\},$$

where

$$\Sigma(T_i) := \bigcup_{g \in G} \bigcup_{j=0}^{\infty} \bigcup_{k=1}^{r_i} g \cdot x_{i,k}^j \cdot g^{-1}.$$

We obtain that the datum of a surface isogenous to a higher product of unmixed type $S = (C_1 \times C_2)/G$ with $q = 0$ is determined, once we look at the monodromy of each covering of $\mathbb{P}^1$, by the datum of a finite group G together with two respective disjoint spherical r_i -systems of generators $T_1 := (x_1, \dots, x_{r_1})$ and $T_2 := (x_1, \dots, x_{r_2})$, such that the types of the systems satisfy (3) with $g' = 0$ and respectively $g = g(C_i)$. The condition of being disjoint ensures that the action of G on the product of the two curves $C_1 \times C_2$ is free. Observe that this can be specialized to $r_i = 3$, and therefore can be used to construct Beauville surfaces. This description suggests the following definition.

Definition 2.7. An unmixed ramification structure of size (r_1, r_2) for a finite group G , is a pair (T_1, T_2) of tuples $T_1 := (x_1, \dots, x_{r_1})$, $T_2 := (x_1, \dots, x_{r_2})$ of elements of G , such that (T_1, T_2) is a disjoint pair of spherical r_i -system of generators of G .

The definition of an *unmixed Beauville structure* given in the introduction is a special case of the above definition for $r_1 = r_2 = 3$. Therefore, the problem of finding Beauville surfaces of unmixed type is now translated into the problem of finding finite groups G which admit an unmixed Beauville structure.

3. RAMIFICATION STRUCTURES FOR FINITE SIMPLE GROUPS

3.1. Ramification Structures for A_n and S_n . In this section we prove Theorems 1.3 and 1.4. The proofs are based on results of Liebeck and Shalev [LS04].

3.1.1. Theoretical Background – Higman’s Conjecture and a Theorem of Liebeck and Shalev on Fuchsian groups. Conder [Co80] (following Higman) proved that sufficiently large alternating groups are in fact *Hurwitz groups*, namely they are quotients of the Hurwitz triangle group $\Delta(2, 3, 7)$, using the method of coset diagrams. In fact, Higman had already conjectured in the late 1960s that every hyperbolic triangle group, and more generally – every Fuchsian group, subjects to all but finitely many alternating groups.

This conjecture was proved by Everitt [Ev] using the method of coset diagrams, and later Liebeck and Shalev [LS04] gave an alternative proof based on probabilistic group theory. In fact, they proved a more explicit and general result, which is presented below.

Note that the results of Liebeck and Shalev are applicable to any Fuchsian group Γ , however, we shall use them only for the case of orbifold surface groups (see Definition 2.4)

$$\Gamma = \Gamma(g' \mid m_1, \dots, m_r)$$

that satisfy the inequality

$$(5) \quad 2g' - 2 + \sum_{i=1}^r \left(1 - \frac{1}{m_i}\right) > 0.$$

Definition 3.1. Let $C_i = g_i^{S_n}$ ($1 \leq i \leq r$) be conjugacy classes in S_n , and let m_i be the order of g_i . Define $\text{sgn}(C_i) = \text{sgn}(g_i)$, and write $\mathbf{C} = (C_1, \dots, C_r)$. Define

$$\text{Hom}_{\mathbf{C}}(\Gamma, S_n) = \{\phi \in \text{Hom}(\Gamma, S_n) : \phi(c_i) \in C_i \text{ for } 1 \leq i \leq r\}.$$

Definition 3.2. Conjugacy classes in S_n of cycle-shape (m^k) , where $n = mk$, namely, containing k cycles of length m each, are called homogeneous. A conjugacy class having cycle-shape $(m^k, 1^f)$, namely, containing k cycles of length m each and f fixed points, with f bounded, is called almost homogeneous.

Theorem 3.3. [LS04, Theorem 1.9]. Let Γ be a Fuchsian group, and let C_i ($1 \leq i \leq r$) be conjugacy classes in S_n with cycle-shapes $(m_i^{k_i}, 1^{f_i})$, where $f_i < f$ for some constant f and $\prod_{i=1}^r \text{sgn}(C_i) = 1$. Set $\mathbf{C} = (C_1, \dots, C_r)$. Then the probability that a random homomorphism in $\text{Hom}_{\mathbf{C}}(\Gamma, S_n)$ has image containing A_n tends to 1 as $n \rightarrow \infty$.

Applying this when Γ is the triangle group $\Delta(m_1, m_2, m_3)$ demonstrates that three elements, with product 1, from almost homogeneous classes C_1, C_2, C_3 of orders m_1, m_2, m_3 , randomly generate A_n or S_n , provided $1/m_1 + 1/m_2 + 1/m_3 < 1$. In particular, when $(m_1, m_2, m_3) = (2, 3, 7)$, this gives random $(2, 3, 7)$ generation of A_n .

Using Theorem 3.3, Liebeck and Shalev deduced the following Corollary regarding symmetric groups.

Corollary 3.4. [LS04, Theorem 1.10]. Let $\Gamma = \Gamma(- \mid m_1, \dots, m_r)$ be a polygonal group which satisfies the above inequality (5), and assume that at least two of $m_1, \dots, m_r$ are even. Then Γ surjects to all but finitely many symmetric groups S_n .

3.1.2. Beauville Structures and Ramification Structures for A_n and S_n .

Proof of Theorem 1.3. Assume that (r_1, s_1, t_1) and (r_2, s_2, t_2) are two hyperbolic types and that n is large enough. By the following Algorithm 3.5, we choose six almost homogeneous conjugacy classes in S_n , $C_{r_1}, C_{s_1}, C_{t_1}, C_{r_2}, C_{s_2}, C_{t_2}$, of orders $r_1, s_1, t_1, r_2, s_2, t_2$ respectively, such that they contain only even permutations, and they all have different numbers of fixed points.

By Theorem 3.3, the probability that three random elements (x_1, y_1, z_1) (equivalently (x_2, y_2, z_2)) whose product is 1, taken from the almost homogeneous conjugacy classes $(C_{r_1}, C_{s_1}, C_{t_1})$ (equivalently $(C_{r_2}, C_{s_2}, C_{t_2})$) will generate A_n , tends to 1 as $n \rightarrow \infty$.

This implies that if n is large enough, one can find six elements $x_1, y_1, z_1, x_2, y_2, z_2$ in A_n of orders $r_1, s_1, t_1, r_2, s_2, t_2$ respectively satisfying the following properties.

- $x_1 \in C_{r_1}, y_1 \in C_{s_1}, z_1 \in C_{t_1}, x_2 \in C_{r_2}, y_2 \in C_{s_2}, z_2 \in C_{t_2}$.
- $x_1 y_1 z_1 = x_2 y_2 z_2 = 1$ and $\langle x_1, y_1 \rangle = \langle x_2, y_2 \rangle = A_n$.
- For any choice of integers $l_{x_1}, l_{y_1}, l_{z_1}, l_{x_2}, l_{y_2}, l_{z_2}$, if the six elements $x_1^{l_{x_1}}, y_1^{l_{y_1}}, z_1^{l_{z_1}}, x_2^{l_{x_2}}, y_2^{l_{y_2}}, z_2^{l_{z_2}}$ are not trivial, then they all belong to different conjugacy classes in S_n , since they all have different numbers of fixed points, and hence $\Sigma(x_1, y_1, z_1) \cap \Sigma(x_2, y_2, z_2) = \{1_{A_n}\}$.

Therefore, if n is large enough, the quadruple $(x_1, y_1; x_2, y_2)$ is an unmixed Beauville structure for A_n , where (x_1, y_1, z_1) has type (r_1, s_1, t_1) and (x_2, y_2, z_2) has type (r_2, s_2, t_2) . $\square$

Algorithm 3.5. *Choosing six almost homogeneous conjugacy classes $C_{r_1}, C_{s_1}, C_{t_1}, C_{r_2}, C_{s_2}, C_{t_2}$ in S_n , of orders $r_1, s_1, t_1, r_2, s_2, t_2$ respectively, such that they contain only even permutations, and they all have different numbers of fixed points.*

Step 1: Sorting $r_1, s_1, t_1, r_2, s_2, t_2$.

Let $m_6 \leq \dots \leq m_1$ be the sorted sequence whose elements are exactly $r_1, s_1, t_1, r_2, s_2, t_2$. Since n can be as large as we want, we may assume that $n > 100m_1$.

Step 2: Choosing even integers k'_i ($1 \leq i \leq 6$).

For $1 \leq i \leq 6$, let

$$k'_i = \begin{cases} \lfloor n/m_i \rfloor & \text{if it is even,} \\ \lfloor n/m_i \rfloor - 1 & \text{otherwise.} \end{cases}$$

Observe that for $1 \leq i \leq 6$,

$$k'_i m_i \leq n \leq (k'_i + 2)m_i.$$

Step 3: Choosing even integers k_i ($1 \leq i \leq 6$) s.t. for every $1 \leq i \neq j \leq 6$, $k_i m_i \neq k_j m_j$.

It may happen that for some $i \neq j$, $k'_i m_i = k'_j m_j$. Therefore, for every i we will choose from the set $\{k'_i - 2l : 0 \leq l \leq 5\}$ a proper integer $k_i = k'_i - 2l$ (for some l), s.t. for every $1 \leq i \neq j \leq 6$, $k_i m_i \neq k_j m_j$. Note that by our assumption, the integers k_i ($1 \leq i \leq 6$) are positive.

Step 4: Defining the conjugacy classes C_i ($1 \leq i \leq 6$).

Assume that n is large enough and let C_i ($1 \leq i \leq 6$) be conjugacy classes in S_n with cycle shapes

$$(m_i^{k_i}, 1^{f_i}), \text{ where } f_i = n - k_i m_i.$$

Observe that the conjugacy classes C_i ($1 \leq i \leq 6$) satisfy the following properties:

- (i) For every $1 \leq i \leq 6$, $\text{sgn}(C_i) = 1$, since C_i contains an even number of cycles (as the k_i -s are even).

- (ii) Set $f := 12m_1$. Then for every $1 \leq i \leq 6$,

$$f_i = n - k_i m_i \leq (k'_i + 2)m_i - (k'_i - 10)m_i = 12m_i \leq 12m_1 = f,$$
and hence it is bounded independently of n .
- (iii) For every $1 \leq i \neq j \leq 6$, $f_i \neq f_j$, since $k_i m_i \neq k_j m_j$.
- (iv) Let $c_i \in C_i$ be some element, then any non-trivial power $c_i^{l_i}$ has exactly f_i fixed points.
- (v) By (iii) and (iv), for any $1 \leq i \neq j \leq 6$ and any two integers l_i, l_j , if the powers $c_i^{l_i}$ and $c_j^{l_j}$ are not trivial, then they belong to different conjugacy classes in S_n .

Step 5: Defining the conjugacy classes $C_{r_1}, C_{s_1}, C_{t_1}, C_{r_2}, C_{s_2}, C_{t_2}$.

Let $k_{r_1}, k_{s_1}, k_{t_1}, k_{r_2}, k_{s_2}, k_{t_2}$ (respectively $f_{r_1}, f_{s_1}, f_{t_1}, f_{r_2}, f_{s_2}, f_{t_2}$) be the elements of the set $\{k_1, \dots, k_6\}$ (respectively $\{f_1, \dots, f_6\}$), ordered by the same correspondence between $\{r_1, s_1, t_1, r_2, s_2, t_2\}$ and $\{m_1, \dots, m_6\}$.

Now, $C_{r_1}, C_{s_1}, C_{t_1}, C_{r_2}, C_{s_2}, C_{t_2}$ are the six conjugacy classes in S_n with cycle-shapes $(r_1^{k_{r_1}}, 1^{f_{r_1}}), (s_1^{k_{s_1}}, 1^{f_{s_1}}), (t_1^{k_{t_1}}, 1^{f_{t_1}}), (r_2^{k_{r_2}}, 1^{f_{r_2}}), (s_2^{k_{s_2}}, 1^{f_{s_2}}), (t_2^{k_{t_2}}, 1^{f_{t_2}})$ respectively.

In a similar way, we prove Theorem 1.4 regarding the symmetric groups.

Proof of Theorem 1.4. Assume that (r_1, s_1, t_1) and (r_2, s_2, t_2) are two hyperbolic types, such that at least two of (r_1, s_1, t_1) are even and at least two of (r_2, s_2, t_2) are even, and that n is large enough. By slightly modifying Algorithm 3.5, we may choose six almost homogeneous conjugacy classes $C_{r_1}, C_{s_1}, C_{t_1}, C_{r_2}, C_{s_2}, C_{t_2}$ in S_n , of orders $r_1, s_1, t_1, r_2, s_2, t_2$ respectively, such that two classes of $C_{r_1}, C_{s_1}, C_{t_1}$ and two classes of $C_{r_2}, C_{s_2}, C_{t_2}$ contain only odd permutations, and all these classes have different numbers of fixed points.

By Theorem 3.3 and Corollary 3.4, the probability that three random elements (x_1, y_1, z_1) (equivalently (x_2, y_2, z_2)) whose product is 1, taken from the almost homogeneous conjugacy classes $(C_{r_1}, C_{s_1}, C_{t_1})$ (equivalently $(C_{r_2}, C_{s_2}, C_{t_2})$) will generate S_n , tends to 1 as $n \rightarrow \infty$.

Therefore, if n is large enough, there exists a quadruple $(x_1, y_1; x_2, y_2)$ which is an unmixed Beauville structure for S_n , where (x_1, y_1, z_1) has type (r_1, s_1, t_1) and (x_2, y_2, z_2) has type (r_2, s_2, t_2) . $\square$

Moreover, since Theorem 3.3 and Corollary 3.4 apply to any polygonal group, one can modify Algorithm 3.5 and deduce the following Corollaries.

Corollary 3.6. Let $\tau_1 = (m_{1,1}, \dots, m_{1,r_1})$ and $\tau_2 = (m_{1,1}, \dots, m_{1,r_2})$ be two sequences of natural numbers such that $m_{k,i} \geq 2$ and $\sum_{i=1}^{r_k} (1 - 1/m_{k,i}) > 2$ for $k = 1, 2$. Then almost all alternating groups A_n admit an unmixed ramification structure of type (τ_1, τ_2) .

Corollary 3.7. Let $\tau_1 = (m_{1,1}, \dots, m_{1,r_1})$ and $\tau_2 = (m_{1,1}, \dots, m_{1,r_2})$ be two sequences of natural numbers such that $m_{k,i} \geq 2$, at least two of $(m_{k,1}, \dots, m_{k,r_k})$ are even and $\sum_{i=1}^{r_k} (1 - 1/m_{k,i}) > 2$, for $k = 1, 2$. Then almost all symmetric groups S_n admit an unmixed ramification structure of type (τ_1, τ_2) .

3.2. Beauville Structures for $\text{PSL}(2, p^e)$. In this section we prove Theorem 1.5 and Corollary 1.6. The proof is based on well-known properties of $\text{PSL}(2, p^e)$ (see for example [Di, Go, Su]) and on results of Macbeath [Ma].

3.2.1. Theoretical Background I – Properties of $\mathrm{PSL}(2, p^e)$. Let $q = p^e$, where p is a prime number and $e \geq 1$. Recall that $\mathrm{GL}(2, q)$ is the group of invertible 2×2 matrices over the finite field with q elements, which we denote by $\mathbb{F}_q$, and $\mathrm{SL}(2, q)$ is the subgroup of $\mathrm{GL}(2, q)$ comprising the matrices with determinant 1. Then $\mathrm{PGL}(2, q)$ and $\mathrm{PSL}(2, q)$ are the quotients of $\mathrm{GL}(2, q)$ and $\mathrm{SL}(2, q)$ by their respective centers.

When q is even, then one can identify $\mathrm{PSL}(2, q)$ with $\mathrm{SL}(2, q)$ and also with $\mathrm{PGL}(2, q)$, and so its order is $q(q-1)(q+1)$. When q is odd, the orders of $\mathrm{PGL}(2, q)$ and $\mathrm{PSL}(2, q)$ are $q(q-1)(q+1)$ and $\frac{1}{2}q(q-1)(q+1)$ respectively, and therefore we can identify $\mathrm{PSL}(2, q)$ with a normal subgroup of index 2 in $\mathrm{PGL}(2, q)$. Also recall that $\mathrm{PSL}(2, q)$ is simple for $q \neq 2, 3$.

One can classify the elements of $\mathrm{PSL}(2, q)$ according to the possible Jordan forms of their pre-images in $\mathrm{SL}(2, q)$. The following table lists the three types of elements, according to whether the characteristic polynomial $P(\lambda) := \lambda^2 - \alpha\lambda + 1$ of the matrix $A \in \mathrm{SL}(2, q)$ (where α is the trace of A) has 0, 1 or 2 distinct roots in $\mathbb{F}_q$.

element type	roots of $P(\lambda)$	canonical form in $\mathrm{SL}(2, \overline{\mathbb{F}}_p)$	order in $\mathrm{PSL}(2, q)$	conjugacy classes
unipotent	1 root	$\begin{pmatrix} \pm 1 & 1 \\ 0 & \pm 1 \end{pmatrix}$ $\alpha = \pm 2$	p	two conjugacy classes in $\mathrm{PSL}(2, q)$, which unite in $\mathrm{PGL}(2, q)$
split	2 roots	$\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix}$ where $a \in \mathbb{F}_q^*$ and $a + a^{-1} = \alpha$	divides $\frac{1}{d}(q-1)$ $d = 1$ for q even $d = 2$ for q odd	for each α : one conjugacy class in $\mathrm{PSL}(2, q)$
non-split	no roots	$\begin{pmatrix} a & 0 \\ 0 & a^q \end{pmatrix}$ where $a \in \mathbb{F}_{q^2}^* \setminus \mathbb{F}_q^*$ $a^{q+1} = 1$ and $a + a^q = \alpha$	divides $\frac{1}{d}(q+1)$ $d = 1$ for q even $d = 2$ for q odd	for each α : one conjugacy class in $\mathrm{PSL}(2, q)$

The subgroups of $\mathrm{PSL}(2, q)$ are well-known (see [Di, Su]), and fall into the following three classes.

Class I: The small triangle subgroups.

These are the *finite* triangle groups $\Delta = \Delta(l, m, n)$, which can occur if and only if $1/l + 1/m + 1/n > 1$.

This inequality holds only for the following triples:

- $(2, 2, n)$: Δ is a dihedral subgroup of order $2n$.
- $(2, 3, 3)$: $\Delta \cong A_4$.
- $(2, 3, 4)$: $\Delta \cong S_4$.
- $(2, 3, 5)$: $\Delta \cong A_5$.

Moreover, if at least two of l, m and n are equal to 2 or if $2 \leq l, m, n \leq 5$, then a subgroup of $\mathrm{PSL}(2, q)$ which is generated by three elements x, y and $z = (xy)^{-1}$, of orders l, m and n respectively, may be a small triangle group (for a detailed list of such triples see [Ma, §8]).

Class II: Structural subgroups.

Let $\mathcal{B}$ be a subgroup of $\mathrm{PSL}(2, q)$ defined by the images of the matrices

$$\left\{ \begin{pmatrix} a & b \\ 0 & a^{-1} \end{pmatrix} : a \in \mathbb{F}_q^*, b \in \mathbb{F}_q \right\},$$

and let $\mathcal{C}$ be a subgroup of $\mathrm{PSL}(2, \overline{\mathbb{F}}_q)$ defined by the images of the matrices

$$\left\{ \begin{pmatrix} t & 0 \\ 0 & t^q \end{pmatrix} : t \in \mathbb{F}_{q^2} \setminus \mathbb{F}_q, t^{q+1} = 1 \right\}.$$

Any subgroup of $\mathrm{PSL}(2, q)$ which can be conjugated (in $\mathrm{PSL}(2, \overline{\mathbb{F}}_q)$) to a subgroup of either $\mathcal{B}$ or $\mathcal{C}$ is called a *structural subgroup* of $\mathrm{PSL}(2, q)$.

Class III: Subfield subgroups.

If $\mathbb{F}_{p^r}$ is a subfield of $\mathbb{F}_q$, then $\mathrm{PSL}(2, p^r)$ is a subgroup of $\mathrm{PSL}(2, q)$. If the quadratic extension $\mathbb{F}_{p^{2r}}$ is also a subfield of $\mathbb{F}_q$, then $\mathrm{PGL}(2, p^r)$ is a subgroup of $\mathrm{PSL}(2, q)$. These groups, as well as any other subgroup of $\mathrm{PSL}(2, q)$ which is isomorphic to any one of them, will be referred to as *subfield subgroups* of $\mathrm{PSL}(2, q)$.

3.2.2. Theoretical Background II – Generation Theorems of Macbeath. Let $(\alpha, \beta, \gamma) \in \mathbb{F}_q^3$, and denote

$$E(\alpha, \beta, \gamma) := \{A, B, C \in \mathrm{SL}(2, q) : ABC = I, \mathrm{tr}A = \alpha, \mathrm{tr}B = \beta, \mathrm{tr}C = \gamma\}.$$

Since all elements in $\mathrm{PSL}(2, q)$ whose pre-images in $\mathrm{SL}(2, q)$ have the same trace are conjugate in $\mathrm{PGL}(2, q)$, all of them have the same order in $\mathrm{PSL}(2, q)$. Therefore, we may denote by $\mathrm{Ord}(\alpha)$ the order in $\mathrm{PSL}(2, q)$ of the image of a matrix $A \in \mathrm{SL}(2, q)$ whose trace equals α .

Example 3.8. $\mathrm{Ord}(0) = 2$, $\mathrm{Ord}(\pm 1) = 3$ and $\mathrm{Ord}(\pm 2) = p$.

Theorem 3.9. [Ma, Theorem 1]. $E(\alpha, \beta, \gamma) \neq \emptyset$ for any $(\alpha, \beta, \gamma) \in \mathbb{F}_q^3$.

Definition 3.10. Let $(\alpha, \beta, \gamma) \in \mathbb{F}_q^3$. We say that (α, β, γ) is *singular* if

$$\alpha^2 + \beta^2 + \gamma^2 - \alpha\beta\gamma = 4.$$

Let $l = \mathrm{Ord}(\alpha)$, $m = \mathrm{Ord}(\beta)$ and $n = \mathrm{Ord}(\gamma)$. We say that (α, β, γ) is *small* if at least two of l, m, n are equal to 2 or if $2 \leq l, m, n \leq 5$.

Theorem 3.11. [Ma, Theorem 2]. $(\alpha, \beta, \gamma) \in \mathbb{F}_q^3$ is *singular* if and only if for $(A, B, C) \in E(\alpha, \beta, \gamma)$, the group generated by the images of A and B is a structural subgroup of $\mathrm{PSL}(2, q)$.

Theorem 3.12. [Ma, Theorem 4]. If $(\alpha, \beta, \gamma) \in \mathbb{F}_q^3$ is *neither singular nor small*, then for any $(A, B, C) \in E(\alpha, \beta, \gamma)$, the group generated by the images of A and B is a subfield subgroup of $\mathrm{PSL}(2, q)$.

Macbeath [Ma] used these generation theorems of $\mathrm{PSL}(2, q)$ to prove that $\mathrm{PSL}(2, q)$ can be generated by two elements one of which is an involution. Moreover, he classified all the values of q for which $\mathrm{PSL}(2, q)$ is a *Hurwitz group*, namely a quotient of the Hurwitz triangle group $\Delta(2, 3, 7)$. In addition, he deduced the following.

Corollary 3.13. [Ma, Theorem 7]. If p is an odd prime, then $\mathrm{PSL}(2, p)$ can be generated by two unipotents whose product is also unipotent.

3.2.3. Beauville Structures for $\mathrm{PSL}(2, p^e)$.

Proof of 1.5. It is known by [BCG05, Proposition 3.6] (and can be easily verified by computer calculations) that $\mathrm{PSL}(2, 2) \cong S_3$, $\mathrm{PSL}(2, 3) \cong A_4$ and $\mathrm{PSL}(2, 4) \cong \mathrm{PSL}(2, 5) \cong A_5$ do not admit an unmixed Beauville structure.

Case $q = p^e$ odd.

Let $q \geq 13$ be an odd prime power, then we will construct an unmixed Beauville structure for $\mathrm{PSL}(2, q)$, $(A_1, B_1; A_2, B_2)$, of type (τ_1, τ_2) , where

$$\tau_1 = \left(\frac{q-1}{2}, \frac{q-1}{2}, \frac{q-1}{2} \right) \text{ and } \tau_2 = \left(\frac{q+1}{2}, \frac{q+1}{2}, \frac{q+1}{2} \right).$$

Let $r = \frac{q-1}{2}$ (respectively $r = \frac{q+1}{2}$), and note that $r > 5$. Let α be a trace of some diagonal split (respectively non-split) element $A \in \mathrm{SL}(2, q)$ whose image in $\mathrm{PSL}(2, q)$ has exact order r , and note that $\alpha \neq 0, \pm 1, \pm 2$, since A is neither of orders 2 or 3 nor unipotent (see Example 3.8).

Observe that (α, α, α) is a non-singular triple. Indeed, the equality $3\alpha^2 - \alpha^3 = 4$ is equivalent to $(\alpha - 2)^2(\alpha + 1) = 0$, but the latter is not possible.

By Theorem 3.9, $E(\alpha, \alpha, \alpha) \neq \emptyset$, and since (α, α, α) is not singular nor small, for $(A, B, C) \in E(\alpha, \alpha, \alpha)$, one has $A \neq \pm B$, and moreover, the image of the subgroup $\langle A, B \rangle$ is a subfield subgroup of $\mathrm{PSL}(2, q)$, by Theorem 3.12. However, since the order of A is exactly $\frac{q-1}{2}$ (respectively $\frac{q+1}{2}$) then the image of the subgroup $\langle A, B \rangle$ is exactly $\mathrm{PSL}(2, q)$.

Observe that $\frac{q-1}{2}$ and $\frac{q+1}{2}$ are relatively prime. Hence, if $A_1, A_2 \in \mathrm{PSL}(2, q)$ have orders $\frac{q-1}{2}$ and $\frac{q+1}{2}$ respectively, then every two non-trivial powers A_1^i and A_2^j have different orders, thus

$$\{g_1 A_1^i g_1^{-1}\}_{g_1, i} \cap \{g_2 A_2^j g_2^{-1}\}_{g_2, j} = \{1\},$$

implying that $\Sigma(A_1, B_1, C_1) \cap \Sigma(A_2, B_2, C_2) = \{1\}$, as needed.

For smaller values of q , a computer calculation (using MAGMA) shows that $\mathrm{PSL}(2, 7)$ admits an unmixed Beauville structure of type $((4, 4, 4), (7, 7, 7))$, $\mathrm{PSL}(2, 9) \cong A_6$ admits an unmixed Beauville structure of type $((4, 4, 4), (5, 5, 5))$, and $\mathrm{PSL}(2, 11)$ admits an unmixed Beauville structure of type $((5, 5, 5), (6, 6, 6))$.

Case $q = 2^e$ even.

Let $q \geq 8$ be an even prime power, then we will construct an unmixed Beauville structure for $\mathrm{PSL}(2, q)$, $(A_1, B_1; A_2, B_2)$, of type (τ_1, τ_2) , where

$$\tau_1 = (q-1, q-1, q-1) \text{ and } \tau_2 = (q+1, q+1, q+1).$$

Let $r = q-1$ (respectively $r = q+1$), and note that $r > 5$. Let α be a trace of some diagonal split (respectively non-split) element $A \in \mathrm{PSL}(2, q) = \mathrm{SL}(2, q)$ of exact order r , and note that $\alpha \neq 0, 1$, since A is neither unipotent nor of order 3 (see Example 3.8).

Observe that (α, α, α) is a non-singular triple. Indeed, the equality $\alpha^2 + \alpha^2 + \alpha^2 - \alpha^3 = 4$ is equivalent (in characteristic 2) to $\alpha^2 + \alpha^3 = \alpha^2(\alpha+1) = 0$, but the latter is not possible.

By Theorem 3.9, $E(\alpha, \alpha, \alpha) \neq \emptyset$, and since (α, α, α) is not singular nor small, for $(A, B, C) \in E(\alpha, \alpha, \alpha)$, one has $A \neq B$, and moreover, the subgroup $\langle A, B \rangle$ is a subfield subgroup of $\mathrm{PSL}(2, q)$, by Theorem 3.12.

However, since the order of A is exactly $q - 1$ (respectively $q + 1$), then $\langle A, B \rangle = \text{PSL}(2, q)$.

Observe that $q - 1$ and $q + 1$ are relatively prime (since both of them are odd). Hence, if $A_1, A_2 \in \text{PSL}(2, q)$ have orders $q - 1$ and $q + 1$ respectively, then every two non-trivial powers A_1^i and A_2^j have different orders, thus

$$\{g_1 A_1^i g_1^{-1}\}_{g_1, i} \cap \{g_2 A_2^j g_2^{-1}\}_{g_2, j} = \{1\},$$

implying that $\Sigma(A_1, B_1, C_1) \cap \Sigma(A_2, B_2, C_2) = \{1\}$, as needed. $\square$

Observe that the above proof actually shows that the group $\text{PSL}(2, q)$ can admit many Beauville structures of various types. On the other hand, if the types are fixed then we can deduce the following.

Corollary 3.14. *Let p be an odd prime, and let $r, s > 5$ be two relatively prime integers which divide the order of $\text{PSL}(2, p)$. Then $\text{PSL}(2, p)$ admits an unmixed Beauville structure of type $((r, r, r), (s, s, s))$.*

Proof. Since r (respectively s) divides $|\text{PSL}(2, p)|$ then it must divide either $\frac{p-1}{2}$ or $\frac{p+1}{2}$ or p . If both r and s divide $\frac{p-1}{2}$ then the result immediately follows from the proof of Theorem 1.5. Otherwise, if $r = p$ then it relies on Corollary 3.13 as well. $\square$

proof of Corollary 1.6. Without loss of generality we may assume that s is odd. By the Chinese Remainder Theorem there exists a unique integer $0 \leq x < 2rs$ solving the system of simultaneous congruences

$$x \equiv 1 \pmod{2r}, \quad x \equiv -1 \pmod{s}.$$

Note that such x necessarily satisfies that $r \mid \frac{x-1}{2}$ and $s \mid \frac{x+1}{2}$.

Moreover, by Dirichlet's Theorem, the arithmetic progression $A_{r,s} := \{2rsn + x : n \in \mathbb{N}\}$ contains infinitely many primes.

By Corollary 3.14 the group $\text{PSL}(2, p)$ admits an unmixed Beauville structure of type $((r, r, r), (s, s, s))$. $\square$

The following two Remarks explain why Theorem 1.3 fails to hold in its great generality for the family of groups $\text{PSL}(2, q)$.

Remark 3.15. Note that unlike the case of Alternating groups, for the group $\text{PSL}(2, p)$, the condition that r and s are relatively prime is also *necessary*.

Indeed, assume that r and s are not relatively prime. Then either $r = s = p$; or both r and s divide either $\frac{p-1}{2}$ or $\frac{p+1}{2}$.

Assume that A_1 and A_2 are two elements in $\text{PSL}(2, p)$ of orders r and s respectively and that both r and s divide $\frac{p-1}{2}$ (or $\frac{p+1}{2}$). Let $d = \gcd(r, s)$, then $r = r'd$ and $s = s'd$ where $\gcd(r', s') = 1$. Now, $A_1^{r'}$ and $A_2^{s'}$ are both of exact order d , hence the cyclic subgroups $\langle A_1^{r'} \rangle$ and $\langle A_2^{s'} \rangle$ are conjugate in $\text{PSL}(2, p)$, implying that there exist some integers k, l such that $A_1^{r'k}$ and $A_2^{s'l}$ are conjugate to the same element of order d .

If A_1 and A_2 are both of order p , then A_i ($i = 1, 2$) can be conjugated in $\text{PSL}(2, p)$ to the image of some matrix $\begin{pmatrix} 1 & a_i \\ 0 & 1 \end{pmatrix}$, where $a_1, a_2 \in \mathbb{F}_p^*$. Since half the elements in $S := \{k : 1 \leq k \leq p-1\}$ are squares in $\mathbb{F}_p^*$ and half are

non-squares, there exist $k, l \in S$ such that A_1^k and A_2^l are both conjugate in $\mathrm{PSL}(2, p)$ to the image of $\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}$.

Remark 3.16. Note that Corollary 3.14 does not hold for the family of groups $\{\mathrm{PSL}(2, p^e)\}_{p \text{ prime}, e \in \mathbb{N}}$, since one cannot fix a hyperbolic type (r, s, t) and hope that almost all groups $G = \mathrm{PSL}(2, p^e)$ where r, s and t all divide $|G|$, will be quotients of $\Delta(r, s, t)$.

Indeed, Macbeath [Ma, Theorem 8] proved that $\mathrm{PSL}(2, p^e)$ is a *Hurwitz group*, namely a quotient of $\Delta(2, 3, 7)$, if either $e = 1$ and $p = 0, \pm 1 \pmod{7}$, or $e = 3$ and $p = \pm 2, \pm 3 \pmod{7}$. Recently, Marion [Mar09] showed that this phenomenon occurs in general for any prime hyperbolic type. Namely, he showed that if (r, s, t) is a hyperbolic triple of primes and p is a prime number, then there exists a unique integer e such that $\mathrm{PSL}(2, p^e)$ is a quotient of the triangle group $\Delta(r, s, t)$.

Interestingly, this situation is different for other families of groups of Lie type of low Lie rank (under the assumption that (r, s, t) are not too small), as was shown in recent results of Marion [Mar3.09, Mar9.09], which are detailed in Theorem 3.17 below.

3.3. Beauville Structures for Other Finite Simple Groups of Lie Type. In this section we prove Theorem 1.7 regarding certain families of finite simple groups of Lie type of low Lie rank. The proof is based on recent results of Marion [Mar3.09, Mar9.09]. Moreover, we discuss some Conjectures on finite simple groups of Lie type in general.

3.3.1. Beauville Structures for Finite Simple Groups of Low Lie Rank.

Theorem 3.17. [Mar3.09, Theorems 1,2,4] and [Mar9.09, Theorem 1]. *Let G be one of the finite simple groups of Lie type listed below, and let (p_1, p_2, p_3) be a hyperbolic tripe of primes $p_1 \leq p_2 \leq p_3$, such that $\mathrm{lcm}(p_1, p_2, p_3)$ divides $|G|$, which, moreover, satisfy the conditions given bellow.*

- (1) *Suzuki groups, $G = {}^2B_2(q)$, where $q = 2^{2e+1}$;*
- (2) *Ree groups, $G = {}^2G_2(q)$, where $q = 3^{2e+1}$;*
- (3) *$G = G_2(q)$, where $q = p^e$ for some prime number $p > 3$, and $(p_1, p_2, p_3) \notin \{(2, 5, 5), (3, 3, 5), (3, 5, 5), (5, 5, 5)\}$;*
- (4) *$G = {}^3D_4(q)$, where $q = p^e$ for some prime number $p > 3$, and (p_1, p_2, p_3) are distinct primes, s.t. $\{p_1, p_2\} \neq \{2, 3\}$;*
- (5) *$G = \mathrm{PSL}(3, q)$, where $q = p^e$ for some prime p , and (p_1, p_2, p_3) are odd primes;*
- (6) *$G = \mathrm{PSU}(3, q)$, where $q = p^e$ for some prime p , and (p_1, p_2, p_3) are odd primes.*

Then, if $\phi \in \mathrm{Hom}(\Delta, G)$ is a randomly chosen homomorphism from the triangle group $\Delta = \Delta(p_1, p_2, p_3)$ to G , then

$$\lim_{q \rightarrow \infty} \mathrm{Prob}\{\phi \text{ is surjective}\} = 1.$$

Now we have all the ingredients needed for the proof of Theorem 1.7.

Proof of Theorem 1.7. (1) Let $G = {}^2B_2(q)$, where $q = 2^{2e+1}$, then

$$|G| = q^2(q^2 + 1)(q - 1).$$

Since $q^2 + 1 \equiv 0 \pmod{5}$, there are at least two prime numbers, 5 and some $r > 5$, which divide $|G|$. If q is large enough, then, by Theorem 3.17, the two triangle groups, $\Delta(5, 5, 5)$ and $\Delta(r, r, r)$, surject onto G , and hence G admits a Beauville structure of type $((5, 5, 5), (r, r, r))$.

(2) Let $G = {}^2G_2(q)$, where $q = 3^{2e+1}$, then

$$|G| = q^3(q-1)(q^3+1).$$

Since $q^3 + 1 \equiv 0 \pmod{7}$, there are at least two odd prime numbers, 7 and some r ($7 \neq r > 3$), which divide $|G|$. If q is large enough, then, by Theorem 3.17, the two triangle groups, $\Delta(7, 7, 7)$ and $\Delta(r, r, r)$, surject onto G , and hence G admits a Beauville structure of type $((7, 7, 7), (r, r, r))$.

(3) Let $G = G_2(q)$, where $q = p^e$ for some prime number $p > 3$, then

$$|G| = q^6(q-1)^2(q+1)^2(q^2-q+1)(q^2+q+1),$$

(which is a factorization into irreducible polynomials), and so there are at least two distinct prime numbers, $r, s \geq 7$, which divide $|G|$. If q is large enough, then, by Theorem 3.17, the two triangle groups, $\Delta(r, r, r)$ and $\Delta(s, s, s)$, surject onto G , and hence G admits a Beauville structure of type $((s, s, s), (r, r, r))$.

(4) Let $G = {}^3D_4(q)$, where $q = p^e$ for some prime number $p > 3$, then

$$|G| = q^{12}(q-1)^2(q+1)^2(q^2-q+1)^2(q^2+q+1)^2(q^4-q^2+1),$$

(which is a factorization into irreducible polynomials), and so there are at least six distinct primes, $p_1 = 2, p_2 = 3, p_3, p_4, p_5, p_6$, which divide $|G|$. If q is large enough, then, by Theorem 3.17, the two triangle groups, $\Delta(2, p_3, p_5)$ and $\Delta(3, p_4, p_6)$, surject onto G , and hence G admits a Beauville structure of type $((2, p_3, p_5), (3, p_4, p_6))$.

(5) Let $G = \text{PSL}(3, q)$ (resp. $G = \text{PSU}(3, q)$), where $q = p^e$ for some prime p , then

$$|G| = \frac{1}{d}q^3(q-1)^2(q+1)(q^2+q+1),$$

(resp. $|G| = \frac{1}{d}q^3(q-1)(q+1)^2(q^2-q+1)$), where $d = 1$ or 3 .

Hence, there are at least two distinct odd prime numbers, greater than 3, r and s , which divide $|G|$. If q is large enough, then, by Theorem 3.17, the two triangle groups, $\Delta(r, r, r)$ and $\Delta(s, s, s)$, surject onto G , and hence G admits a Beauville structure of type $((s, s, s), (r, r, r))$. $\square$

3.3.2. Conjectures on Finite Simple Classical Groups of Lie Type. Liebeck and Shalev raised the following Conjecture in [LS05] regarding finite simple classical groups of Lie type.

Conjecture 3.18 (Liebeck-Shalev). For any Fuchsian group Γ there is an integer $f(\Gamma)$, such that if G is a finite simple classical group of Lie rank at least $f(\Gamma)$, then the probability that a randomly chosen homomorphism from Γ to G is an epimorphism tends to 1 as $|G| \rightarrow \infty$.

If this Conjecture holds, it immediately implies that any finite simple classical group G of Lie rank large enough admits an unmixed Beauville structure. Indeed, let s and t be two distinct primes greater than 3, then the triangle groups $\Delta(s, s, s)$ and $\Delta(t, t, t)$ will surject onto G , if G is of Lie

rank large enough, yielding a Beauville structure of type $((s, s, s), (t, t, t))$ for G . Moreover, this Conjecture inspired us to formulate Conjecture 1.8.

Acknowledgement. The authors are grateful to Fritz Grunewald for inspiring and motivating us to work on this problem together. He is deeply missed.

The authors would like to thank Ingrid Bauer and Fabrizio Catanese for suggesting the problems, for many useful discussions and for their helpful suggestions, especially concerning the Theorem of Liebeck and Shalev.

We are grateful to Bob Guralnick, Martin Liebeck, Alex Lubotzky, Martin Kassabov, Aner Shalev, Boris Kunyavskii and Eugene Plotkin for interesting discussions. We would like to thank Claude Marion for kindly referring us to his recent results.

The authors acknowledge the support of the DFG Forschergruppe 790 "Classification of algebraic surfaces and compact complex manifolds". The first author acknowledges the support of the European Post-Doctoral Institute and the Max-Planck-Institute for Mathematics in Bonn.

REFERENCES

- [BHPV] W.P. Barth, K. Hulek, C.A.M. Peters, A. Van de Ven, *Compact complex surfaces*. Second edition. Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics], **4**. Springer-Verlag, Berlin, 2004.
- [BCG05] I. Bauer, F. Catanese, F. Grunewald, *Beauville surfaces without real structures*. In: Geometric methods in algebra and number theory, Progr. Math., vol **235**, Birkhäuser Boston, (2005) 1–42.
- [BCG06] I. Bauer, F. Catanese, F. Grunewald, *Chebycheff and Belyi polynomials, dessins d'enfants, Beauville surfaces and group theory*. Mediterr. J. Math. **3**, no.2, (2006) 121–146.
- [BCG08] I. Bauer, F. Catanese, F. Grunewald, *The classification of surfaces with $p_g = q = 0$ isogenous to a product*. Pure Appl. Math. Q., **4**, no. 2, part1, (2008) 547–586.
- [B] A. Beauville, *Surfaces Algébriques Complex*. Astérisque **54**, Paris (1978).
- [Cat00] F. Catanese, *Fibred surfaces, varieties isogenous to a product and related moduli spaces*. Amer. J. Math. **122**, (2000) 1–44.
- [Co80] M.D.E. Conder, *Generators for alternating and symmetric groups*. J. London Math. Soc. **22** (1980) 75–86.
- [Co90] M.D.E. Conder, *Hurwitz groups: a brief survey*. Bull. Amer. Math. Soc. **23** (1990) 359–370.
- [Co10] M.D.E. Marston, *An update on Hurwitz groups*. Groups, Complexity and Cryptology **2**, no. 1, (2010) 35–49.
- [Di] L. E. Dickson. *Linear groups with an exposition of the Galois field theory*. (Teubner, 1901).
- [Ev] B. Everitt, *Alternating quotients of Fuchsian groups*. J. Algebra **223** (2000) 457–476.
- [FG] Y. Fuertes, G. González-Diez, *On Beauville structures on the groups S_n and A_n* , Math. Z. **264**, no. 4, (2010) 959–968.
- [FMP] B. Fairbairn, K. Magaard, C. Parker, *Generation of finite simple groups with an application to groups acting on Beauville surfaces*, Preprint available at arXiv:1010.3500.
- [FGJ] Y. Fuertes, G. González-Diez, A. Jaikin-Zapirain, *On Beauville surfaces*. Preprint.
- [FJ] Y. Fuertes, G. Jones, *Beauville surfaces and finite groups*. Preprint available at arXiv:0910.5489.
- [GLL] S. Garion, M. Larsen, A. Lubotzky, *Beauville surfaces and finite simple groups*, Preprint available at arXiv:1005.2316.
- [Go] D. Gorenstein, *Finite groups*. Chelsea Publishing Co., New York, 1980.

- [GM] R. Guralnick, G. Malle, *Simple groups admit Beauville structures*, Preprint available at arXiv:1009.6183.
- [LS04] M.W. Liebeck, A. Shalev, *Fuchsian groups, coverings of Riemann surfaces, subgroup growth, random quotients and random walks*. J. Algebra **276** (2004) 552–601.
- [LS05] M.W. Liebeck, A. Shalev, *Fuchsian groups, finite simple groups and representation varieties*. Invent. Math. **159** (2005) no. 2, 317–367.
- [Ma] A. M. Macbeath, *Generators of the linear fractional groups*, Number Theory (Proc. Sympos. Pure Math., Vol. XII, Houston, Tex., 1967). Amer. Math. Soc., Providence, R.I. (1969) 14–32.
- [Mar09] C. Marion, *Triangle groups and $\mathrm{PSL}_2(q)$* , J. Group Theory **12** (2009) 689–708
- [Mar3.09] C. Marion, *Triangle generation of finite exceptional groups of low rank*. Preprint.
- [Mar9.09] C. Marion, *Random and deterministic triangle generation of three-dimensional classical groups I*. Preprint.
- [S] F. Serrano, *Isotrivial fibred surfaces*. Annali di Matematica. pura e applicata, Vol. **CLXXI**, (1996) 63–81.
- [Su] M. Suzuki, *Group Theory I*. Springer-Verlag, Berlin, 1982.

SHELLY GARION, MAX-PLANCK-INSTITUTE FOR MATHEMATICS, D-53111 BONN, GERMANY

E-mail address: `shellyg@mpim-bonn.mpg.de`

MATTEO PENEGINI, LEHRSTUHL MATHEMATIK VIII, UNIVERSITÄT BAYREUTH, NWII, D-95440 BAYREUTH, GERMANY

E-mail address: `matteo.penegini@uni-bayreuth.de`